# INDEX


Alfalfa, breeding for self-tripping character to improve seed-yield, 82–8; methods of selecting, 82–3; seed-yields, 82–7; yields of single crosses, 83–6, of synthetic strains, 86–7.
Armstrong, J. M., and Stevenson, T. M., effects of continuous line selection in Triticum-Agropyron hybrids, 51–64.

Blue tick, the arsenic-resistant, in South Africa, 42–7.
Bolton, J. L., *see* Stevenson, T. M.
Boyns, B. M., Sudanese cattle as milk-producers, 27–41.
Broadfoot, Jane, preliminary experiments on the protection of cut sets of potato from infection with Penicillium, 227–36.
Bruin, P., *see* Vries, O. de.

Cattle, artificial insemination in, 149–59; breeding policy, 155–8; organization in England, 150–5; technical developments and research, 158–9.
— Indian, in the United States of America, 213–15; hybrids of Brahman (Zebu), English, and native cattle, 213–15; influence of Zebu blood on range cattle, 215; Palmetta palm and other herbage, 214–15.
— infestations with arsenic-resistant blue tick, 42–50.
— pedigree, age-distribution of, 119–31; age in relation to milk-recording and tuberculin-testing, &c., 126–28; age-composition of Red Poll female population, 120–4; age-distribution of Ayrshire and Jersey cows, 125–6; comparison of age-distributions from herd-lists and heifer registrations, 124–5, replacements, 128–30.
— Sudanese, as milk-producers, 27–41; cattle of the Sudan, 27–8; early work on improvement, 28–9; foundation of Shambat herd, 29–30; lactation results, 30–4; methods for improving native stock, 34–40.
Coffee, arabica, yields, preparation out-turns, and quality in. I. Yields, 96–106; biennial bearing, 101–2; climate, soil, seed, 96–7; combined seasons' yields, 103–5; comparison of estate- and plot-yields, 98–9; effect of shade on yields, 102–3; methods used in crop-recording, plot-selection, &c., 97–8; variation in yields, 99–101. II. Preparation out-turns and quality, 167–76; methods, 167–9; quality, 172–4; variation in grades, 171, in out-turn ratios, 169–71, in size and weight of beans, 171–2, in quality, 172–4.
— robusta, in Uganda, 65–81; breeding, 74–9; development, 66–8; historical, 65–6; prospects, 79–80; selection, 68–73; training, 73–4.
Confounding arrangements, construction of, 107–12; factors at two levels, 107–9, three levels, 109–11, more than three levels, 111–12.

Davis, A. G., lucerne growing, 113–18.
Dent, J. M., soil problems of empoldered rice lands in Sierra Leone, 206–12.
Diseases and pests, ecto-parasitic infestations of farm stock in South Africa, 42–50.
Donald, H. P., age-distribution of pedigree cattle and factors affecting it, 119–31.

Edwards, J., artificial insemination in cattle, 149–59.
El Itriby, A. A., *see* Donald, H. P.

Fertilizers, inorganic, compared with compost on South African pastures, 189–94; harvesting methods, 190–1; recovery of N and $P_2O_5$, 192–3; treatments, 189–90; types of pasture, 190; yields, 191–2.
Finney, D. J., construction of confounding arrangements, 107–12.
Foodstuffs, mineral composition of mine-compound, in Northern Rhodesia, 89–95; experimental, 89–90; results, 90–4.

'Gammexane', use of, against arsenic-resistant blue tick in South Africa, 42–50; dipping tests and dip-washes, 44–5; spraying test, 43; trials, object and results of, 45–7.
Glover, J., environment and the growth of the potato in tropical East Africa, 9–26.
Golledge, A., *see* Young, R. S.

Hallsworth, E. G., factors affecting the ascorbic-acid content of tomatoes in New South Wales, 132–47.
Hawkes, J. G., photoperiodic reactions of potato bolters, 216–26.

Insemination, artificial, in cattle, 149–59.

Lathbury, R. J., development of wheat varieties in Kenya, 177–88.
Lewis, V. M., *see* Hallsworth, E. G.
Liming problems on Netherland light soils, 260–8; lime factor, determination of, 260–1; lime losses by leaching and drainage, 262–8; seasonal variations in soil pH, 261–2.
Littlejohn, L. J. S., experiments with potatoes in Cyprus, 195–205.
Lucerne growing, 113–18; companion plants and seed-rates, 113–17; management, 117; 'strain', 117.

Mann, H. H., pulse grain crops in the Middle East, 249–59.
Meredith, D., comparison of inorganic fertilizers and compost on natural pastures in South Africa, 189–94.
Moorland plants, ash components of, 237–48; experimental technique, 237–8; *Eriophorum vaginatum*, 244–5, 246; *Juncus squarrosus*, 243–4; *Molinia coerulea*, 238–9; *Nardus stricta*, 242–3; *Scirpus caespitosus*, 239–40; *Vaccinium myrtillus*, 240–2.

Plank, J. E. van der, some climatic factors determining high yields of potatoes, 1–8.
Potato bolters, photoperiodic reactions of, 216–26; materials and methods, 217; observations on flowering, 218–19; height of plants, 217–18, maturity, 219, stolon length and number, 222–4; tuber length and number, 219–21; summaries, 224, 225–6.
— environment and growth in tropical East Africa, 9–26; environmental factors, 11–17; experimental plots and technique, 9–11; field observations, 18–20; growth and yield measurements, 17.
— protection of cut sets from infection with Penicillium, 227–36; laboratory tests, 228–31; methods used, 227–8; planting trials, 231–2; transport trials, 232–4.
Potatoes, climatic factors determining high yields of. II. At low latitudes and high altitudes, 1–8; length of day, 3–5; poor seed, 2; unsuitable varieties, 2–3. Summary of Pts. I and II, 7–8.
— experiments with, in Cyprus, 195–205; dormancy, breaking of, 202–4; rose-ends, trials with, 198–202; seed certification, 204–5; variety trials, 196–8; yield as function of age of seed, 195.
Pulse grain crops in the Middle East, 249–59; beans, 254–6; chickpeas, 250–1; cowpeas, 256–7; *Lathyrus ochrus*, 258; *Lathyrus sativus*, 257–8; lentils, 253–4; lupins, 258; peas, 256; vetches, 251–3.

Rice lands, soil problems of empoldered, in Sierra Leone, 206–12; soil analyses, 210; toxicity associated with common salt, 211, iron salts, 207 sqq., mangrove and other vegetation, 208–12, peaty matter, 208 sqq.

Stevenson, T. M., evaluation of the self-tripping character in breeding for improved seed-yield in alfalfa, 82–8.

Tea, effect of cultivation on growth of young, 160–6; cultivation methods, 160–2; growth of plants, 162–3, of roots, 162–5, 166.
Thomas, A. S., cultivation and selection of robusta coffee in Uganda, 65–81.
Thomas, Brynmor; ash components of some moorland plants, 237–49.
Thorburn, J. A., control of ecto-parasitic infestations of farm stock with 'Gammexane', with special reference to the arsenic-resistant blue tick, 42–50.
Thorold, C. A., study of yields, preparation out-turns, and quality in arabica coffee, 96–106, 167–76.
Tomatoes, ascorbic-acid content of, in New South Wales, 132–47; climatic data, 132–3; effects of maturity of bush, 138–9, ripeness, 134, storage, 134; relation between content and leafiness and surface area, 136, 142–4, locality, 141–2, size and weight of fruit, times of picking and sowing, 139, 140; varietal differences, 140–4.
Trinder, N., *see* Thomas, Brynmor.
Triticum–Agropyron hybrids, effects of continuous line-selection in, 51–64; chemical composition of hybrids, 54–6; emergence of grasses, T–A hybrids, and wheat sown at varying depth, 53–4; perennialism, 56–8; variability in advanced generations, 58–63.
Tubbs, F. R., effect of cultivation on growth of young tea, 160–6.

Vries, O. de, liming problems on light soils in the Netherlands, 260–8.

Ware, F., Indian cattle in the United States of America, 213–15.
Wheat, development of varieties in Kenya, 177–88; acreage, prices, yields, 178–9, 187; baking quality, 179; breeding for rust-resistance, 181; early history, 177; new hybrids, 181–5; selections and introduced varieties, 185–6; stem rust and yellow rust, 179–80.

Young, R. S., mineral composition of mine-compound foodstuffs in Northern Rhodesia, 89–95.

## CORRIGENDA

Vol. XV, No. 57, pp. 9–26. Fig. 1, *delete* present title and *insert* 'The climate of M'.
— Fig. 2, *for* 'K 1' *read* 'K 2'. *Delete* '21° C.', and after '24° C.' *add* '27° C. o–o and 30° C. x–x'.
— Fig. 3, *delete* present title and *insert* 'The climate of K 2'.
— Fig. 5, *delete* present title and *insert* 'The climate of K 1'.
— Fig. 6, *for* 'K 2' *read* 'K 1'; *delete* '27° C. o–o' and 30° C. x–x', and *insert* '21° C. and' before '24° C.'.